arXiv:1606.05983v2 [math.DG] 7 Feb 2017

COMPLEX AND LAGRANGIAN SURFACES OF THE COMPLEX PROJECTIVE PLANE VIA KÄHLERIAN KILLING SPIN^c SPINORS

ROGER NAKAD AND JULIEN ROTH

ABSTRACT. The complex projective space $\mathbb{C}P^2$ of complex dimension 2 has a Spin^c structure carrying Kählerian Killing spinors. The restriction of one of these Kählerian Killing spinors to a surface M^2 characterizes the isometric immersion of M^2 into $\mathbb{C}P^2$ if the immersion is either Lagrangian or complex.

1. INTRODUCTION

A classical problem in Riemannian geometry is to know when a Riemannian manifold (M^n, g) can be isometrically immersed into a fixed Riemannian manifold $(\widetilde{M}^{n+p}, \widetilde{g})$. The case of space forms $\mathbb{R}^{n+1}$, $\mathbb{S}^{n+1}$ and $\mathbb{H}^{n+1}$ is well-known. In fact, the Gauss, Codazzi and Ricci equations are necessary and sufficient conditions. In other ambient spaces, the Gauss, Codazzi and Ricci equations are necessary but not sufficient in general. Some additional conditions may be required like for the case of complex space forms, products, warped products or 3-dimensional homogeneous spaces (see [8, 9, 21, 24, 33, 36]).

In low dimensions, especially for surfaces, another necessary and sufficient condition is now well-known, namely the existence of a special spinor field called *generalized Killing spinor field* ([10, 29, 23, 25]). These results are the geometrical invariant versions of previous works on the spinorial Weierstrass representation by R. Kusner and N. Schmidt, B. Konoplechenko, I. Taimanov and many others (see [22, 20, 38]). This representation was expressed by T. Friedrich [10] for surfaces in $\mathbb{R}^3$ and then extended to other 3-dimensonal (pseudo-)Riemannian manifolds [29, 36, 35, 26] as well as for hypersurfaces of 4-dimensional space forms and products [25] or hypersurfaces of 2-dimensional complex space forms by means of Spin^c spinors [32].

More precisely, the restriction φ of a parallel spinor field on $\mathbb{R}^{n+1}$ to an oriented Riemannian hypersurface M^n is a solution of the generalized Killing equation

$$\nabla_X \varphi = -\frac{1}{2} A(X) \cdot \varphi, \tag{1}$$

where " $\cdot$ " and ∇ are respectively the Clifford multiplication and the spin connection on M^n, the tensor A is the Weingarten tensor of the immersion and X any vector field on M. Conversely, T. Friedrich proved in [10] that, in the two dimensional case, if there exists a generalized Killing spinor field satisfying Equation (1), where A is an arbitrary field of symmetric endomorphisms of TM, then A satisfies the fundamental Codazzi and Gauss equations in the theory of embedded hypersurfaces in a Euclidean space and consequently, A is the Weingarten tensor of a local isometric immersion of M into $\mathbb{R}^3$. Moreover, in this case, the solution φ of the generalized Killing equation is equivalently a solution of the Dirac equation

$$D\varphi = H\varphi, \tag{2}$$

2010 *Mathematics Subject Classification.* 53C27, 53C40, 53D12, 53C25.

Key words and phrases. Spin^c structures, Kählerian Killing Spin^c spinors, isometric immersions, complex and Lagrangian surfaces, the Dirac operator.

where D denotes the Dirac operator on M, $|\varphi|$ is constant and H is a real-valued function (which is the mean curvature of the immersion in $\mathbb{R}^3$).

More recently, this approach was adapted by the second author, P. Bayard and M.A. Lawn in codimension two, namely, for surfaces in Riemannian 4-dimensional real space forms [4], and then generalized in the pseudo-Riemannian setting [3, 5] as well as for 4-dimensional products [37]. As pointed out in [34], this approach coincides with the Weierstrass type representation for surfaces in $\mathbb{R}^4$ introduced by Konopelchenko and Taimanov [20, 39].

The aim of the present article is to provide an analogue for the complex projective space $\mathbb{C}P^2$. The key point is that, contrary to the case of hypersurfaces of $\mathbb{C}P^2$ which was considered in [32], the use of Spin^c parallel spinors is not sufficient. Indeed, for both canonical or anti-canonical Spin^c structures, the parallel spinors are always in the positive half-part of the spinor bundle. But, as proved in [4] or [37], a spinor with non-vanishing positive and negative parts is required to get the integrability condition of an immersion in the desired target space. For this reason, Spin^c parallel spinors are not adapted to our problem. Thus, we make use of real Kählerian Killing spinors. Therefore, our argument holds for the complex projective space and not for the complex hyperbolic space, since $\mathbb{C}\mathbb{H}^2$ does not carry a real or imaginary Kählerian Killing spinor.

We will focus on the case of complex and Lagrangian immersions into $\mathbb{C}P^2$. These two cases and especially the Lagrangian case are of particular interest in the study of surfaces in $\mathbb{C}P^2$ (see [14, 15, 40] and references therein for instance).

First, consider (M^2, g) an oriented Riemannian Spin^c surface and E an oriented Spin^c vector bundle of rank 2 over M with scalar product $\langle\cdot,\cdot\rangle_E$ and compatible connection ∇^E. We denote by F^M (resp. F^E) the curvature form (an imaginary 2-form on M) of the auxiliary line bundle defining the Spin^c structure on M (resp. on the vector bundle E). For a spinor field φ, we define $\bar\varphi$ by $\bar\varphi = \varphi_+ - \varphi_-$, where φ^+ and φ^- denote the positive and negative half part of φ (see Section 2). They are the projections of φ on the eigensubspaces for the eigenvalues $+1$ and -1 of the complex volume form. The aim of the paper is to prove the following two results:

The first theorem gives a spinorial characterization of complex immersions of surfaces in the complex projective space $\mathbb{C}P^2$.

Theorem 1.1. *Let (M^2, g) be an oriented Riemannian surface and E an oriented vector bundle of rank 2 over M with scalar product $\langle\cdot,\cdot\rangle_E$ and compatible connection ∇^E. We denote by $\Sigma = \Sigma M \otimes \Sigma E$ the twisted spinor bundle. Let $B : TM \times TM \longrightarrow E$ be a bilinear symmetric map, $j : TM \longrightarrow TM$ a complex structure on M and $t : E \longrightarrow E$ a complex structure on E. Assume moreover that $t(B(X,Y)) = B(X, j(Y))$ for all $X \in \Gamma(TM)$ and consider $\{e_1, e_2\}$ an orthonormal frame of TM. Then, the following two statements are equivalent*

(1) *There exists a Spin^c structure on $\Sigma M \otimes \Sigma E$ whose auxiliary line bundle's curvature is given by $F^{M+E}(e_1, e_2) := F^M(e_1, e_2) + F^E(e_1, e_2) = 0$ and a spinor field $\varphi \in \Gamma(\Sigma M \otimes \Sigma E)$ satisfying for all $X \in \Gamma(TM)$,*

$$\nabla_X \varphi \quad = \quad -\frac{1}{2}\eta(X)\cdot\varphi - \frac{1}{2}X\cdot\varphi + \frac{i}{2}j(X)\cdot\overline{\varphi}, \tag{3}$$

such that φ^+ and φ^- never vanish and where η is given by

$$\eta(X) = \sum_{j=1}^{2} e_j \cdot B(e_j, X).$$

(2) *There exists a local isometric* **complex** *immersion of (M^2, g) into $\mathbb{C}P^2$ with E as normal bundle and second fundamental form B such that the complex structure of $\mathbb{C}P^2$ over M is given by j and t (in the sense of Proposition 3.2).*

The second theorem is the analogue of Theorem 1.1 for Lagrangian surfaces in $\mathbb{C}P^2$.

Theorem 1.2. *Let (M^2, g) be an oriented Riemannian surface and E an oriented vector bundle of rank 2 over M with scalar product $\langle\cdot,\cdot\rangle_E$ and compatible connection ∇^E. We denote by $\Sigma = \Sigma M \otimes \Sigma E$ the twisted spinor bundle. Let $B : TM \times TM \longrightarrow E$ be a bilinear symmetric map, $h : TM \longrightarrow E$ a bundle map and $s : E \longrightarrow TM$ the dual map of h. Assume moreover that h and s are parallel, $h \circ s = -\mathrm{id}_E$ and $A_{h(Y)}X + s(B(X,Y)) = 0$, for all $X \in \Gamma(TM)$, where $A_\nu : TM \longrightarrow TM$ is defined by $g(A_\nu X, Y) = \langle B(X,Y), \nu\rangle_E$ for all $X, Y \in \Gamma(TM)$ and $\nu \in E$. Then, the following two statements are equivalent*

(1) *There exists a Spin^c structure on $\Sigma M \otimes \Sigma E$ whose auxiliary line bundle's curvature is given by $F^{M+E}(e_1, e_2) = -2i$ and a spinor field $\varphi \in \Gamma(\Sigma M \otimes \Sigma E)$ satisfying for all $X \in \Gamma(M)$,*

$$\nabla_X \varphi \quad = \quad -\frac{1}{2}\eta(X)\cdot\varphi - \frac{1}{2}X\cdot\varphi + \frac{i}{2}h(X)\cdot\overline{\varphi}, \tag{4}$$

such that φ^+ and φ^- never vanish and where η is given by

$$\eta(X) = \sum_{j=1}^{2} e_j \cdot B(e_j, X).$$

(2) *There exists a local isometric* **Lagrangian** *immersion of (M^2, g) into $\mathbb{C}P^2$ with E as normal bundle and second fundamental form B such that over M the complex structure of $\mathbb{C}P^2$ is given by h and s (in the sense of Proposition 3.2).*

2. Preliminaries and Notation

In this section, we briefly review some basic facts about Kähler geometry and Spin^c structures on manifolds and their submanifolds. For more details we refer to [30, 2, 6, 27, 28, 16, 17, 7, 1, 11, 12, 13].

2.1. **Spin^c structures on Kähler-Einstein manifolds.** Let (M^n, g) be an n-dimensional closed Riemannian Spin^c manifold and denote by ΣM its complex spinor bundle, which has complex rank equal to $2^{[\frac{n}{2}]}$. The bundle ΣM is endowed with a Clifford multiplication denoted by "$\cdot$" and a scalar product denoted by $\langle\cdot,\cdot\rangle$. Given a Spin^c structure on (M^n, g), one can check that the determinant line bundle $\det(\Sigma M)$ has a root L of index $2^{[\frac{n}{2}]-1}$. This line bundle L over M is called the auxiliary line bundle associated with the Spin^c structure. In the particular case, when the auxiliary line bundle can be chosen to be trivial, the manifold is called a spin manifold. The connection ∇ on ΣM is the twisted connection of the one on the spin bundle (induced by the Levi-Civita connection) and a fixed connection A on L. The Spin^c Dirac operator D acting on the space of sections of ΣM is defined locally by $D = \sum_{j=1}^{n} e_j \cdot \nabla_{e_j}$, where $\{e_j\}_{j=1,\ldots,n}$ is a local orthonormal basis of TM.

We recall that the complex volume element $\omega_{\mathbb{C}} = i^{[\frac{n+1}{2}]} e_1 \wedge \ldots \wedge e_n$ acts as the identity on the spinor bundle if n is odd. If n is even, $\omega_{\mathbb{C}}^2 = 1$. Thus, under the action of the complex volume element, the spinor bundle decomposes into the eigenspaces $\Sigma^{\pm}M$ corresponding to the ± 1 eigenspaces, the *positive* and the *negative* spinors.

Every Kähler manifold (M^{2m}, g, J) has a canonical Spin^c structure induced by the complex structure J. The complexified tangent bundle decomposes into $T^{\mathbb{C}}M = T_{1,0}M \oplus T_{0,1}M$, the i-eigenbundle (resp. $(-i)$-eigenbundle) of the complex linear extension of J. For any vector field X, we denote by $X^{\pm} := \frac{1}{2}(X \mp iJ(X))$ its component in $T_{1,0}M$, resp. $T_{0,1}M$. The spinor bundle of the canonical Spin^c structure is defined by

$$\Sigma M = \Lambda^{0,*}M = \bigoplus_{r=0}^{m} \Lambda^r(T_{0,1}^* M),$$

and its auxiliary line bundle is $L = (\mathcal{K}_M)^{-1} = \Lambda^m(T^*_{0,1}M)$, where $\mathcal{K}_M = \Lambda^{m,0}M$ is the canonical bundle of M. The line bundle L has a canonical holomorphic connection, whose curvature form is given by $-i\rho$, where ρ is the Ricci form defined, for all vector fields X and Y, by $\rho(X,Y) = \mathrm{Ric}(J(X),Y)$ and Ric denotes the Ricci tensor. Similarly, one defines the so called anti-canonical Spin^c structure, whose spinor bundle is given by $\Lambda^{*,0}M = \oplus_{r=0}^m \Lambda^r(T^*_{1,0}M)$ and the auxiliary line bundle by $\mathcal{K}_M$. The spinor bundle of any other Spin^c structure on M can be written as:

$$\Sigma M = \Lambda^{0,*}M \otimes \mathbb{L},$$

where $\mathbb{L}^2 = \mathcal{K}_M \otimes L$ and L is the auxiliary line bundle associated with this Spin^c structure. The Kähler form Ω, defined as $\Omega(X,Y) = g(J(X),Y)$, acts on ΣM via Clifford multiplication. Under this action, the spinor bundle decomposes as follows:

$$\Sigma M = \bigoplus_{r=0}^{m} \Sigma_r M, \tag{5}$$

where $\Sigma_r M$ denotes the eigenbundle to the eigenvalue $i(2r-m)$ of Ω, of complex rank $\binom{m}{k}$. It is easy to see that $\Sigma_r M \subset \Sigma^+ M$ (resp. $\Sigma_r M \subset \Sigma^- M$) if and only if r is even (resp. r is odd). Moreover, for any $X \in \Gamma(TM)$ and $\varphi \in \Gamma(\Sigma_r M)$, we have $X^+ \cdot \varphi \in \Gamma(\Sigma_{r+1}M)$ and $X^- \cdot \varphi \in \Gamma(\Sigma_{r-1}M)$, with the convention $\Sigma_{-1}\mathrm{M} = \Sigma_{m+1}\mathrm{M} = M \times \{0\}$. Thus, for any Spin^c structure, we have $\Sigma_r M = \Lambda^{0,r}M \otimes \Sigma_0 M$. Hence, $(\Sigma_0 M)^2 = \mathcal{K}_M \otimes L$, where L is the auxiliary line bundle associated with the Spin^c structure. For example, when the manifold is spin, we have $(\Sigma_0 M)^2 = \mathcal{K}_M$ [18, 19]. For the canonical Spin^c structure, since $L = (\mathcal{K}_M)^{-1}$, it follows that $\Sigma_0 M$ is trivial. This yields the existence of parallel spinors (the constant functions) lying in $\Sigma_0 M$, *cf.* [31].

In [16], the authors gave examples of Spin^c structures on compact Kähler-Einstein manifolds of positive scalar curvature, which carry Kählerian Killing Spin^c spinors lying in $\Sigma_r M \oplus \Sigma_{r+1}M$, for $r \neq \frac{m\pm1}{2}$, in contrast to the spin case, where Kählerian Killing spinors may only exist for m odd in the middle of the decomposition (5). We briefly describe these Spin^c structures here. If the first Chern class $c_1(\mathcal{K}_M)$ of the canonical bundle of the Kähler manifold M is a non-zero cohomology class, the greatest number $p \in \mathbb{N}^*$ such that $\frac{1}{p}c_1(\mathcal{K}_M) \in H^2(M,\mathbb{Z})$, is called the *Maslov index* of the Kähler manifold. One can thus consider a p-th root of the canonical bundle $\mathcal{K}_M$, *i.e.* a complex line bundle $\mathcal{L}$, such that $\mathcal{L}^p = \mathcal{K}_M$. In [16], O. Hijazi, S. Montiel and F. Urbano proved the following:

Theorem 2.1 (Theorem 14, [16])**.** *Let M be a $2m$-dimensional Kähler-Einstein compact manifold with scalar curvature $4m(m+1)$ and Maslov index index $p \in \mathbb{N}^*$. For each $0 \leq r \leq m+1$, there exists on M a Spin^c structure with auxiliary line bundle given by $\mathcal{L}^q$, where $q = \frac{p}{m+1}(2r-m-1) \in \mathbb{Z}$, and carrying a Kählerian Killing spinor $\psi_{r-1} + \psi_r \in \Gamma(\Sigma_{r-1}M \oplus \Sigma_r M)$,* i.e. *for all $X \in \Gamma(TM)$, it satisfies the first order system*

$$\begin{cases} \nabla_X \psi_r = -X^+ \cdot \psi_{r-1}, \\ \nabla_X \psi_{r-1} = -X^- \cdot \psi_r. \end{cases}$$

For example, if M is the complex projective space $\mathbb{C}P^m$ of complex dimension m and constant holomorphic sectional curvature 4, then $p = m+1$ and $\mathcal{L}$ is just the tautological line bundle. We fix $0 \leq r \leq m+1$ and we endow $\mathbb{C}P^m$ with the Spin^c structure whose auxiliary line bundle is given by $\mathcal{L}^q$ where $q = \frac{p}{m+1}(2r-m-1) = 2r-m-1 \in \mathbb{Z}$. For this Spin^c structure, the space of Kählerian Killing spinors in $\Gamma(\Sigma_{r-1}M \oplus \Sigma_r M)$ has dimension $\binom{m+1}{r}$. In this example, for $r = 0$ (resp. $r = m+1$), we get the canonical (resp. anticanonical) Spin^c structure for which Kählerian Killing spinors are just parallel spinors.

2.2. **Submanifolds of Spin^c manifolds.** Let (M^2, g) be an oriented Riemannian surface, with a given Spin^c structure, and E an oriented Spin^c vector bundle (see [27, Chapter II]) of rank 2 on M with an Hermitian product $\langle\cdot,\cdot\rangle_E$ and a compatible connection ∇^E.

We consider the spinor bundle Σ over M twisted by E and defined by $\Sigma = \Sigma M \otimes \Sigma E$, where ΣM and ΣE are the spinor bundles of M and E respectively. We endow Σ with the spinorial connection ∇ defined by

$$\nabla = \nabla^{\Sigma M} \otimes Id_{\Sigma E} + Id_{\Sigma M} \otimes \nabla^{\Sigma E},$$

where $\nabla^{\Sigma M}$ and $\nabla^{\Sigma E}$ are respectively the Spin^c connections on ΣM and ΣE. We also define the Clifford product "$\cdot$" by

$$\begin{cases} X \cdot \varphi = (X \cdot_{_M} \alpha) \otimes \overline{\sigma} & \text{if } X \in \Gamma(TM), \\ X \cdot \varphi = \alpha \otimes (X \cdot_{_E} \sigma) & \text{if } X \in \Gamma(E), \end{cases}$$

for all $\varphi = \alpha \otimes \sigma \in \Gamma(\Sigma M \otimes \Sigma E)$, where "$\cdot_{_M}$" and "$\cdot_{_E}$" denote the Clifford multiplications on ΣM and on ΣE respectively and where $\overline{\sigma} = \sigma^+ - \sigma^-$ under the natural decomposition of $\Sigma E = \Sigma^+ E \oplus \Sigma^- E$. Here, $\Sigma^+ E$ and $\Sigma^- E$ are the eigensubbundles (for the eigenvalue 1 and -1) of ΣE for the action of the normal volume element $\omega_\perp = i\nu_1 \cdot_E \nu_2$, where $\{\nu_1, \nu_2\}$ is a local orthonormal frame of E. Note that $\Sigma^+ M$ and $\Sigma^- M$ are defined similarly by the action of the tangent volume element $\omega = ie_1 \cdot_M e_2$, where $\{e_1, e_2\}$ is an orthonormal basis of TM. The twisted Dirac operator D on $\Gamma(\Sigma)$ is defined by

$$D\varphi = e_1 \cdot \nabla_{e_1} \varphi + e_2 \cdot \nabla_{e_2} \varphi.$$

We note that Σ is also naturally equipped with a hermitian scalar product $\langle ., . \rangle$ which is compatible with the connection ∇, and thus also with a compatible real scalar product $\mathrm{Re}\langle ., . \rangle$. We also note that the Clifford product "$\cdot$" of vectors belonging to $TM \oplus E$ is antihermitian with respect to this hermitian product. Finally, we stress that the four subbundles $\Sigma^{\pm\pm} := \Sigma^\pm M \otimes \Sigma^\pm E$ are orthogonal with respect to the hermitian product. We will also consider $\Sigma^+ = \Sigma^{++} \oplus \Sigma^{--}$ and $\Sigma^- = \Sigma^{+-} \oplus \Sigma^{-+}$. Throughout the paper we will assume that the hermitian product is $\mathbb{C}-$linear w.r.t. the first entry, and $\mathbb{C}-$antilinear w.r.t. the second entry.

Let $(\widetilde{M}^4, \widetilde{g})$ be a Riemannian Spin^c manifold and (M^2, g) an oriented surface isometrically immersed into $\widetilde{M}$. We denote by NM the normal bundle of M into $\widetilde{M}$. As M is an oriented surface, it is also Spin^c. We denote by $F^{\widetilde{M}}$ (resp. F^M) the curvature 2-form of the auxiliary line bundle $L^{\widetilde{M}}$ (resp. L) associated with the Spin^c structure on $\widetilde{M}$ (resp. M). Since the manifolds M and $\widetilde{M}$ are Spin^c, there exists a Spin^c structure on the bundle NM whose auxiliary line bundle L_N is given by $L_N := (L)^{-1} \otimes L^{\widetilde{M}}{}_{|_M}$. We denote by ΣN the Spin^c bundle of NM and let $\Sigma = \Sigma M \otimes \Sigma N$ the spinor bundle over M twisted by NM constructed as above with the associated connection and Clifford multiplication. It is a classical fact that the spinor bundle of $\widetilde{M}$ over M, $\Sigma\widetilde{M}_{|M}$ identifies with Σ. Moreover the connections on each bundle are linked by the spinorial Gauss formula: for any $\varphi \in \Gamma(\Sigma)$ and any $X \in \Gamma(TM)$,

$$\widetilde{\nabla}_X \varphi = \nabla_X \varphi + \frac{1}{2} \sum_{j=1,2} e_j \cdot B(X, e_j) \cdot \varphi \tag{6}$$

where B is the second fundamental form, $\widetilde{\nabla}$ is the spinorial connection of $\Sigma\widetilde{M}$ and ∇ is the spinoral connection of Σ defined as above and $\{e_1, e_2\}$ is a local orthonormal frame of TM. Here "$\cdot$" is the Clifford product on $\Sigma\widetilde{M}$ which identifies with the Clifford mulitplication on Σ.

3. Immersed surfaces into the complex projective space

In this section, we will give the basic facts about immersed surfaces in the complex projective plane and in particular derive a sequence of necessary and sufficient conditions for the existence of such immersions.

3.1. **Compatibility equations.** Let (M^2, g) be a Riemannian surface isometrically immersed in the 2-dimensional complex projective space of constant holomorphic sectional curvature $4c > 0$. We denote by ∇ the Levi-Civita connection of $\Big(M^2, g := \langle ., . \rangle\Big)$, $\widetilde{g}$ $\Big($also denoted by $\langle ., . \rangle$ without ambiguity$\Big)$ the Fubini-Study metric of $\mathbb{C}P^2(4c)$ and $\widetilde{\nabla}$ its Levi-Civita connection. Moreover, we denote by $\nabla^\perp$ the normal connection and $R^\perp$ the normal curvature. First of all, we recall that for any $X, Y, Z \in \Gamma(TM)$ the curvature tensor of $\mathbb{C}P^2(4c)$ is given by

$$\begin{aligned}\widetilde{R}(X,Y,Z,W) &= c\Big[\langle X,W\rangle\langle Y,Z\rangle - \langle X,Z\rangle\langle Y,W\rangle + \langle J(X),W\rangle\langle J(Y),Z\rangle \\ &\quad - \langle J(X),Z\rangle\langle J(Y),W\rangle + 2\langle X,J(Y)\rangle\langle J(Z),W\rangle\Big]. \qquad (7)\end{aligned}$$

The complex structure J induces the existence of the following four operators

$$j : TM \longrightarrow TM,\ h : TM \longrightarrow NM,\ s : NM \longrightarrow TM \text{ and } t : NM \longrightarrow NM$$

defined for any $X \in \Gamma(TM)$ and $\xi \in \Gamma(NM)$ by

$$J(X) = j(X) + h(X) \quad \text{and} \quad J(\xi) = s(\xi) + t(\xi). \qquad (8)$$

From the fact that $J^2 = -\mathrm{Id}$ and J is antisymmetric, we get that j and t are antisymmetric and for any $X \in \Gamma(TM)$ and $\xi \in \Gamma(NM)$, we have

$$j^2(X) = -X - s \circ h(X), \qquad (9)$$
$$t^2(\xi) = -\xi - h \circ s(\xi), \qquad (10)$$
$$j \circ s(\xi) + s \circ t(\xi) = 0, \qquad (11)$$
$$h \circ j(X) + t \circ h(X) = 0, \qquad (12)$$
$$\langle h(X), \xi\rangle = -\langle X, s(\xi)\rangle. \qquad (13)$$

We denote by $B : TM \times TM \longrightarrow NM$ the second fundamental form and by S_ξ the Weingarten operator associated with $\xi \in \Gamma(NM)$ and defined by $\widetilde{g}(S_\xi X, Y) = \widetilde{g}(B(X,Y), \xi)$ for any vectors X, Y tangent to M. From the fact that J is parallel, we have

$$(\nabla_X j)Y = S_{h(Y)}X + s(B(X,Y)), \qquad (14)$$
$$\nabla^\perp_X(h(Y)) - h(\nabla_X Y) = t(B(X,Y)) - B(X, j(Y)), \qquad (15)$$
$$\nabla^\perp(t(\xi)) - t(\nabla^\perp_X \xi) = -B(s(\xi), X) - h(S_\xi X), \qquad (16)$$
$$\nabla_X(s(\xi)) - s(\nabla^\perp_X \xi) = -j(S_\xi X) + S_{t(\xi)}X, \qquad (17)$$

where Finally, from (7), we deduce that the Gauss, Codazzi and Ricci equations are respectively given by

$$\begin{aligned}R(X,Y)Z &= c\Big[\langle Y,Z\rangle X - \langle X,Z\rangle Y + \langle j(Y),Z\rangle j(X) - \langle j(X),Z\rangle j(Y) \\ &\quad + 2\langle X, j(Y)\rangle j(Z)\Big] + S_{B(Y,Z)}X - S_{B(X,Z)}Y,\end{aligned}$$

$$(\nabla_X B)(Y,Z) - (\nabla_Y B)(X,Z) = c\Big[\langle j(Y),Z\rangle h(X) - \langle j(X),Z\rangle h(Y) + 2\langle j(X),Y\rangle h(Z)\Big],$$

$$\begin{aligned}R^\perp(X,Y)\xi &= c\Big[\langle h(Y),\xi\rangle h(X) - \langle h(X),\xi\rangle h(Y) + 2\langle j(X),Y\rangle t(\xi)\Big] \\ &\quad + B(S_\xi Y, X) - B(S_\xi X, Y).\end{aligned}$$

In the local orthonormal frames $\{e_1, e_2\}$ and $\{\nu_1, \nu_2\}$ and for any $k, l \in \{1, 2\}$, we set $B_{kl} = B(e_k, e_l)$, $j_{kl} = g(j(e_k), e_l)$, $t_{kl} = \widetilde{g}(t(\nu_k), e_l)$ and $h_{kl} = \widetilde{g}(h(e_k), \nu_l)$. Thus, these last equations become:

$$K_M = c + < B_{22}, B_{11} > -|B_{12}|^2 + 3c\,(j_{12})^2, \tag{18}$$

$$K_N = - < [S_{\nu_1}, S_{\nu_2}](e_1), e_2 > + c\Big(h_{21}h_{12} - h_{11}h_{22} + 2j_{12}t_{12}\Big), \tag{19}$$

$$(\nabla_{e_1} B)(e_2, e_k) - (\nabla_{e_2} B)(e_1, e_k) = c\Big(j_{2k}h(e_1) - j_{1k}h(e_2) + 2j_{12}h(e_k)\Big). \tag{20}$$

We want to point out that B_{kl} is a normal vector whereas j_{kl}, t_{kl} and h_{kl} are real numbers. It is clear that Equations (9) to (20) are necessary conditions for surfaces in $\mathbb{C}P^2(4c)$. Conversely, given (M^2, g) a Riemannian surface, E a 2-dimensional vector bundle over M endowed with a scalar product $\overline{g}$ and a compatible connection ∇^E. Let $j : TM \longrightarrow TM$, $h : TM \longrightarrow E$, $s : E \longrightarrow TM$ and $t : E \longrightarrow E$ be four tensors. Note that the metric $\widetilde{g} := \langle ., . \rangle$ is defined on $TM \oplus E$ by

$$\begin{cases} \widetilde{g}(X, Y) = g(X, Y) \quad \text{for any } X, Y \in \Gamma(TM), \\ \widetilde{g}(\nu, \xi) = \overline{g}(\xi, \nu) \quad \text{for any } \nu, \xi \in \Gamma(E), \\ \widetilde{g}(X, \nu) = 0 \quad \text{for any } X \in \Gamma(TM) \text{ and } \nu \in \Gamma(E). \end{cases}$$

Definition 3.1. *We say that $(M, g, E, \overline{g}, \nabla^E, B, j, h, s, t)$ satisfies the compatibility equations for $\mathbb{C}P^2(4c)$ if j and t are antisymmetric, the Gauss, the Codazzi and Ricci equations* (18) (19) (20) *and equations* (9)-(17) *are fulfilled.*

Now, we can state the following classical *Fundamental Theorem* for surfaces of $\mathbb{C}P^2$, which can be found for instance as a special case of the general result of P. Piccione and D. Tausk [33, Theorem 8.1 and Example 8.2].

Proposition 3.2. *If $(M, g, E, \overline{g}, \nabla^E, B, j, h, s, t)$ satisfies the compatibility equations for $\mathbb{C}P^2(4c)$ then, there exists an isometric immersion $\Phi : M \longrightarrow \mathbb{C}P^2(4c)$ such that the normal bundle of M for this immersion is isomorphic to E and such that the second fundamental form II and the normal connection $\nabla^\perp$ are given by B and ∇^E. Precisely, there exists a vector bundle isometry $\widetilde{\Phi} : E \longrightarrow NM$ so that*

$$II = \widetilde{\Phi} \circ B,$$
$$\nabla^\perp \widetilde{\Phi} = \widetilde{\Phi} \nabla^E.$$

Moreover, we have

$$J(\Phi_*(X)) = \Phi_*(j(X)) + \widetilde{\Phi}(h(X)),$$
$$J(\widetilde{\Phi}(\xi)) = \Phi_*(s(X)) + \widetilde{\Phi}(t(\xi)),$$

where J is the canonical complex structure of $\mathbb{C}P^2(4c)$ and this isometric immersion is unique up to an isometry of $\mathbb{C}P^2(4c)$.

3.2. **Special cases.** Two special cases are of particular interest and have been widely studied, the complex and Lagrangian surfaces.

A surface (M^2, g) of $\mathbb{C}P^2(4c)$ is said **complex** if the tangent bundle of M is stable by the complex structure of $\mathbb{C}P^2(4c)$, that is, $J(TM) = TM$. Note that we have necessarily $J(NM) = NM$. Hence, in that case, with the above notations, we have $h = 0$, $s = 0$ and so j and t are respectively almost complex structures on TM and NM. The compatibility equations for complex surfaces of $\mathbb{C}P^2(4c)$ become

$$\begin{cases} h = 0,\ s = 0,\ j^2 = -\mathrm{id}_{TM},\ t^2 = -\mathrm{id}_E \\ \nabla j = 0,\ \nabla^\perp t = 0 \\ t(B(X, Y)) - B(X, j(Y)) = 0,\ \forall X, Y \in \Gamma(TM) \\ K_M = 4c + < B_{22}, B_{11} > -|B_{12}|^2 \\ K_N = - < [S_{\nu_1}, S_{\nu_2}](e_1), e_2 > + 2c \\ (\nabla_{e_1} B)(e_2, e_k) - (\nabla_{e_2} B)(e_1, e_k) = 0 \end{cases} \tag{21}$$

A surface (M^2, g) of $\mathbb{C}P^2(4c)$ is said totally real if $J(TM)$ is transversal to TM. In the particular case when $J(TM) = NM$, we say that (M^2, g) is **Lagrangian**. In that case, we have $j = 0$ and $t = 0$. Hence, the compatibility equations for Lagrangian surfaces of $\mathbb{C}P^2(4c)$ are

$$\begin{cases} j = 0,\ t = 0,\ s \circ h = -\mathrm{id}_{TM},\ h \circ s = -\mathrm{id}_E \\ \nabla s = 0,\ \nabla^\perp h = 0 \\ A_{h(Y)}X + s(B(X,Y)) = 0,\ \forall X, Y \in TM \\ K_M = c + < B_{22}, B_{11} > -|B_{12}|^2 \\ K_N = - < [S_{\nu_1}, S_{\nu_2}](e_1), e_2 > + c(h_{21}h_{12} - h_{11}h_{22}) \\ (\nabla_{e_1} B)(e_2, e_k) - (\nabla_{e_2} B)(e_1, e_k) = 0 \end{cases} \tag{22}$$

4. Restriction of a Kählerian Killing Spin^c spinor and the curvature computation

We consider a special Spin^c structure on $\mathbb{C}P^2$ of constant holomorphic sectional curvature $4c = 4$ carrying a (real) Kählerian Killing spinor φ. For example, on can take $q = -1$ and hence $r = 1$. For this structure, the curvature of the auxiliary line bundle is given by $F^{\mathbb{C}P^2}(X, Y) = -2ig(J(X), Y)$. There exists a spinor $\varphi = \varphi_0 + \varphi_1 \in \Gamma(\Sigma_0\mathbb{C}P^2 \oplus \Sigma_1\mathbb{C}P^2)$ satisfies the following:

$$\begin{cases} \widetilde{\nabla}_X \varphi_0 = -X^- \cdot \varphi_1, \\ \widetilde{\nabla}_X \varphi_1 = -X^+ \cdot \varphi_0, \end{cases}$$

Thus, we have

$$\widetilde{\nabla}_X \varphi = -\frac{1}{2} X \cdot \varphi + \frac{i}{2} J(X) \cdot \overline{\varphi},$$

where $\overline{\varphi} = \varphi_0 - \varphi_1$ is the conjugate of φ for the action of the complex volume element $\omega_4^{\mathbb{C}} = -e_1 \wedge e_2 \wedge e_3 \wedge e_4$. Indeed, $\Sigma_0\mathbb{C}P^2 \subset \Sigma^+\mathbb{C}P^2$ and $\Sigma_1\mathbb{C}P^2 = \Sigma^-\mathbb{C}P^2$. Note also that such as spinor is of constant norm and each part φ_0 and φ_1 does not have any zeros. Indeed, for instance, if φ_0 vanishes at one point, then it must vanish everywhere (as it is obtained by parallel transport) and φ_1 is as a parallel spinor which is not the case for this Spin^c structure.

Now, let M be a surface of $\mathbb{C}P^2$ with normal bundle denoted by NM. By the identification of the Clifford multiplications and the Spin^c Gauss formula, we have

$$\nabla_X \varphi = -\frac{1}{2}\eta(X) \cdot \varphi - \frac{1}{2} X \cdot \varphi + \frac{i}{2} J(X) \cdot \overline{\varphi}.$$

In intrinsic terms, it can be written as

$$\nabla_X \varphi \quad = \quad -\frac{1}{2}\eta(X) \cdot \varphi - \frac{1}{2} X \cdot \varphi + \frac{i}{2} j(X) \cdot \overline{\varphi} + \frac{i}{2} h(X) \cdot \overline{\varphi}, \tag{23}$$

where η is given by

$$\eta(X) = \sum_{j=1}^{2} e_j \cdot B(e_j, X). \tag{24}$$

Here B is the second fundamental form of the immersion, and the operators j and h are those introduced in Section 3. After projection on Σ^+ and Σ^- respectively, we deduce immediately that

$$\begin{cases} \nabla_X \varphi^+ = -\frac{1}{2}\eta(X) \cdot \varphi^+ - \frac{1}{2} X \cdot \varphi^- - \frac{i}{2} j(X) \cdot \varphi^- - \frac{i}{2} h(X) \cdot \varphi^-, \\ \\ \nabla_X \varphi^- = -\frac{1}{2}\eta(X) \cdot \varphi^- - \frac{1}{2} X \cdot \varphi^+ + \frac{i}{2} j(X) \cdot \varphi + + - \frac{i}{2} h(X) \cdot \varphi^+. \end{cases}$$

From this, since $\overline{\varphi} = \varphi^+ - \varphi^-$, we get

$$\nabla_X \overline{\varphi} = -\frac{1}{2}\eta(X)\cdot\overline{\varphi} + \frac{1}{2}X\cdot\overline{\varphi} - \frac{i}{2}j(X)\cdot\varphi - \frac{i}{2}h(X)\cdot\varphi. \tag{25}$$

Now, let us go back to an instrinsic setting by considering (M^2, g) an oriented Riemannian surface and E an oriented vector bundle of rank 2 over M with scalar product $\langle\cdot,\cdot\rangle_E$ and compatible connection ∇^E. We denote by $\Sigma = \Sigma M \otimes \Sigma E$ the twisted spinor bundle. Let $B : TM \times TM \longrightarrow E$ a bilinear symmetric map and $j : TM \longrightarrow TM,\ h : TM \longrightarrow E$ two tensors. We assume that the spinor field $\varphi \in \Gamma(\Sigma)$ satisfies Equation (23). We will compute the spinorial curvature for this spinor field φ in the local orthonormal frames $\{e_1, e_2\}$ of TM and $\{\nu_1, \nu_2\}$ of E. For a sake of simplicity, we can assume that $\{e_1, e_2\}$ is normal at the point $p \in M$ so that at p, $\nabla e_1 = 0$, $\nabla e_2 = 0$ and so $[e_1, e_2] = 0$. Hence, at the point p, we have

$$\begin{aligned}
\nabla_{e_1}\nabla_{e_2}\varphi &= -\frac{1}{2}\nabla_{e_1}(\eta(e_2))\cdot\varphi + \frac{1}{4}\eta(e_2)\cdot\eta(e_1)\cdot\varphi + \frac{1}{4}\eta(e_2)\cdot e_1\cdot\varphi \\
&\quad -\frac{i}{4}\eta(e_2)\cdot j(e_1)\cdot\overline{\varphi} - \frac{i}{4}\eta(e_2)\cdot h(e_1)\cdot\overline{\varphi} + \frac{1}{4}e_2\cdot\eta(e_1)\cdot\varphi \\
&\quad +\frac{1}{4}e_2\cdot e_1\cdot\varphi - \frac{i}{4}e_2\cdot j(e_1)\cdot\overline{\varphi} - \frac{i}{4}e_2\cdot h(e_1)\cdot\overline{\varphi} \\
&\quad +\frac{i}{2}\nabla_{e_1}(j(e_2))\cdot\overline{\varphi} - \frac{i}{4}j(e_2)\cdot\eta(e_1)\cdot\overline{\varphi} + \frac{i}{4}j(e_2)\cdot e_1\cdot\overline{\varphi} \\
&\quad +\frac{1}{4}j(e_2)\cdot j(e_1)\cdot\varphi + \frac{1}{4}j(e_2)\cdot h(e_1)\cdot\varphi + \frac{i}{2}\nabla^{\perp}_{e_1}(h(e_2))\cdot\overline{\varphi} \\
&\quad -\frac{i}{4}h(e_2)\cdot\eta(e_1)\cdot\overline{\varphi} + \frac{i}{4}h(e_2)\cdot e_1\cdot\overline{\varphi} + \frac{1}{4}h(e_2)\cdot j(e_1)\cdot\varphi \\
&\quad +\frac{1}{4}h(e_2)\cdot h(e_1)\cdot\varphi \\
&= \underbrace{-\frac{1}{2}\nabla_{e_1}(\eta(e_2))\cdot\varphi + \frac{1}{4}\eta(e_2)\cdot\eta(e_1)\cdot\varphi + \frac{1}{4}e_2\cdot e_1\cdot\varphi}_{\mathcal{A}_1^{12}} \\
&\quad \underbrace{+\frac{1}{4}\eta(e_2)\cdot e_1\cdot\varphi + \frac{1}{4}e_2\cdot\eta(e_1)\cdot\varphi}_{\mathcal{A}_2^{12}} \\
&\quad \underbrace{+\frac{i}{2}\nabla_{e_1}(j(e_2))\cdot\overline{\varphi} + \frac{i}{2}\nabla^{\perp}_{e_1}(h(e_2))\cdot\overline{\varphi}}_{\mathcal{A}_3^{12}} \\
&\quad \underbrace{-\frac{i}{4}e_2\cdot j(e_1)\cdot\overline{\varphi} + \frac{i}{4}j(e_2)\cdot e_1\cdot\overline{\varphi}}_{\mathcal{A}_4^{12}} \\
&\quad \underbrace{-\frac{i}{4}e_2\cdot h(e_1)\cdot\overline{\varphi} + \frac{i}{4}h(e_2)\cdot e_1\cdot\overline{\varphi}}_{\mathcal{A}_5^{12}} \\
&\quad \underbrace{+\frac{1}{4}j(e_2)\cdot j(e_1)\cdot\varphi}_{\mathcal{A}_6^{12}} + \underbrace{\frac{1}{4}h(e_2)\cdot h(e_1)\cdot\varphi}_{\mathcal{A}_7^{12}} \\
&\quad \underbrace{+\frac{1}{4}\big(j(e_2)\cdot h(e_1)\cdot\varphi + h(e_2)\cdot j(e_1)\cdot\varphi\big)}_{\mathcal{A}_8^{12}}
\end{aligned}$$

$$\underbrace{-\frac{i}{4}\big(\eta(e_2)\cdot h(e_1)\cdot\overline{\varphi}+h(e_2)\cdot\eta(e_1)\cdot\overline{\varphi}\big)}_{\mathcal{A}_9^{12}}$$
$$\underbrace{-\frac{i}{4}\big(\eta(e_2)\cdot j(e_1)\cdot\overline{\varphi}+j(e_2)\cdot\eta(e_1)\cdot\overline{\varphi}\big)}_{\mathcal{A}_{10}^{12}}.$$

We point out that since $[e_1,e_2]=0$, we have $\nabla_{[e_1,e_2]}\varphi=0$. and some terms are vanishing as shown in the following lemma.

Lemma 4.1. *We denote by $\mathcal{A}_1^{21}$ (resp. $\mathcal{A}_2^{21},\cdots,\mathcal{A}_{10}^{21}$) the expression $\mathcal{A}_1^{12}$ (resp. $\mathcal{A}_2^{12},\cdots,\mathcal{A}_{10}^{12}$) when e_1 and e_2 are interchanged. We have*

(1)
$$\mathcal{A}_2^{12}-\mathcal{A}_2^{21}=0 \tag{26}$$

(2)
$$\mathcal{A}_5^{12}-\mathcal{A}_5^{21}=0 \tag{27}$$

(3)
$$\mathcal{A}_3^{12}+\mathcal{A}_9^{12}+\mathcal{A}_{10}^{12}-\mathcal{A}_3^{21}-\mathcal{A}_9^{21}-\mathcal{A}_{10}^{21}=0 \tag{28}$$

(4)
$$\mathcal{A}_6^{12}-\mathcal{A}_6^{21}=-\frac{1}{2}(j_{21})^2e_1\cdot e_2\cdot\varphi \tag{29}$$

(5)
$$\mathcal{A}_7^{12}-\mathcal{A}_7^{21}=\frac{1}{2}[h_{21}h_{12}-h_{11}h_{22}]\nu_1\cdot\nu_2\cdot\varphi \tag{30}$$

(6)
$$\mathcal{A}_4^{12}-\mathcal{A}_4^{21}=ij_{12}\overline{\varphi} \tag{31}$$

(7)
$$\begin{aligned}\mathcal{A}_8^{12}-\mathcal{A}_8^{21}= \ & \tfrac{1}{2}\ \Big(j_{21}h_{11}e_1\cdot\nu_1+j_{21}h_{12}e_1\cdot\nu_2\\ & +j_{21}h_{21}e_2\cdot\nu_1+j_{12}h_{22}e_2\cdot\nu_2\Big)\cdot\varphi\end{aligned} \tag{32}$$

(8)
$$\begin{aligned}&\mathcal{A}_1^{12}-\mathcal{A}_1^{21}\\ =\ & -\frac{1}{2}\sum_{j=1}^{2}e_j\cdot\big((\nabla'_{e_1}B)(e_2,e_j))-(\nabla'_{e_2}B)(e_1,e_j)\big)\cdot\varphi\\ & +\frac{1}{2}g([S_{\nu_1},S_{\nu_2}])(e_1),e_2)\nu_1\cdot\nu_2\cdot\varphi\\ & +\frac{1}{2}\big(|B_{12}|^2-\langle B_{11},B_{22}\rangle\big)e_1\cdot e_2-\frac{1}{2}e_1\cdot e_2\cdot\varphi,\end{aligned} \tag{33}$$

where ∇' is the natural connection on $T^*M\otimes T^*M\otimes E$.

Proof:

(1) Using the definition of η, we get $-\frac{1}{2}B(e_j,X)=e_j\cdot\eta(X)-\eta(X)\cdot e_j$. Hence
$$\mathcal{A}_2^{12}-\mathcal{A}_2^{21}=-\frac{1}{8}B_{21}+\frac{1}{8}B_{12}=0.$$

(2)

$$\begin{aligned}\mathcal{A}_5^{12}-\mathcal{A}_5^{21} &= -\frac{i}{4}(e_2\cdot h(e_1)-h(e_2)\cdot e_1)\cdot\overline{\varphi}+\frac{i}{4}(e_1\cdot h(e_2)-h(e_1)\cdot e_2)\cdot\overline{\varphi}\\ &= \frac{i}{4}(2h_{12}-2h_{21})\cdot\overline{\varphi}=0, \qquad (34)\end{aligned}$$

because X and $h(X)$ are orthogonal for any $X\in\Gamma(TM)$ with respect to the metric $\widetilde{g}$.

(3) First we have

$$\begin{aligned}&\mathcal{A}_3^{12}-\mathcal{A}_5^{21}\\ =\;& \frac{i}{2}\nabla_{e_1}(j(e_2))\cdot\overline{\varphi}+\frac{i}{2}\nabla^{\perp}_{e_1}(h(e_2))\cdot\overline{\varphi}-\frac{i}{2}\nabla_{e_2}(j(e_1))\cdot\overline{\varphi}-\frac{i}{2}\nabla^{\perp}_{e_2}(h(e_1))\cdot\overline{\varphi}\\ =\;& \frac{i}{2}\Big((\nabla_{e_1}j)e_2\cdot\overline{\varphi}+(\nabla_{e_1}h)e_2\cdot\overline{\varphi}-(\nabla_{e_2}j)e_1\cdot\overline{\varphi}-(\nabla_{e_2}h)e_1\cdot\overline{\varphi}\Big)\\ =\;& \frac{i}{2}\Big(s(B_{12})\cdot\overline{\varphi}+S_{h(e_2)}e_1\cdot\overline{\varphi}+t(B_{12})\cdot\overline{\varphi}-B(e_1,j(e_2))\cdot\overline{\varphi}\\ &-s(B_{12})\cdot\overline{\varphi}-S_{h(e_1)}e_2\cdot\overline{\varphi}-t(B_{12})\cdot\overline{\varphi}+B(e_2,j(e_1))\cdot\overline{\varphi}\Big)\\ =\;& \frac{i}{2}\Big(S_{h(e_2)}e_1\cdot\overline{\varphi}-S_{h(e_1)}e_2\cdot\overline{\varphi}-B(e_1,j(e_2))\cdot\overline{\varphi}+B(e_2,j(e_1))\cdot\overline{\varphi}\Big) \qquad (35)\end{aligned}$$

Moreover, we calculate

$$\begin{aligned}-B(e_1,j(e_2))\cdot\overline{\varphi}+B(e_2,j(e_1))\cdot\overline{\varphi} &= -j_{21}B_{11}\cdot\overline{\varphi}+j_{12}B_{22}\cdot\overline{\varphi}\\ &= 2g(j(e_1),e_2)H\cdot\overline{\varphi}=2j_{12}H\cdot\overline{\varphi} \qquad (36)\end{aligned}$$

and

$$\begin{aligned}&S_{h(e_2)}e_1\cdot\overline{\varphi}-S_{h(e_1)}e_2\cdot\overline{\varphi}\\ =\;& -<S_{h(e_1)}e_2,e_1>e_1\cdot\overline{\varphi}-<S_{h(e_1)}e_2,e_2>e_2\cdot\overline{\varphi}\\ &+<S_{h(e_2)}e_1,e_1>e_1\cdot\overline{\varphi}+<S_{h(e_2)}e_1,e_2>e_2\cdot\overline{\varphi}\\ =\;& -<B_{21},h(e_1)>e_1\cdot\overline{\varphi}-<B_{22},h(e_1)>e_2\cdot\overline{\varphi}\\ &+<B_{11},h(e_2)>e_1\cdot\overline{\varphi}+<B_{12},h(e_2)>e_2\cdot\overline{\varphi} \qquad (37)\end{aligned}$$

In addition we have

$$\begin{aligned}&\mathcal{A}_9^{12}-\mathcal{A}_9^{21}\\ =\;& \frac{i}{4}\Big(-e_1\cdot B_{12}\cdot h(e_1)-e_2\cdot B_{22}\cdot h(e_1)-h(e_2)\cdot e_1\cdot B_{11}-h(e_2)\cdot e_2\cdot B_{12}\\ &+e_1\cdot B_{11}\cdot h(e_2)+e_2\cdot B_{12}\cdot h(e_2)+h(e_1)\cdot e_1\cdot B_{12}+h(e_1)\cdot e_2\cdot B_{22}\Big)\cdot\overline{\varphi}\\ =\;& \frac{i}{4}\Big(2<B_{12},h(e_1)>e_1+2<B_{22},h(e_1)>e_2\\ &-2<B_{12},h(e_2)>e_2-2<B_{11},h(e_2)>e_1\Big)\cdot\overline{\varphi} \qquad (38)\end{aligned}$$

and

$$\begin{aligned}&\mathcal{A}_{10}^{12}-\mathcal{A}_{10}^{21}\\ =\;& -\frac{i}{4}\eta(e_2)\cdot j(e_1)\cdot\overline{\varphi}-\frac{i}{4}j(e_2)\cdot\eta(e_1)\cdot\overline{\varphi}+\frac{i}{4}\eta(e_1)\cdot j(e_2)\cdot\overline{\varphi}+\frac{i}{4}j(e_1)\cdot\eta(e_2)\cdot\overline{\varphi}\\ =\;& \frac{i}{4}\Big(e_1\cdot j(e_1)\cdot B_{11}+e_2\cdot j(e_1)\cdot B_{22}-j(e_2)\cdot e_1\cdot B_{11}-j(e_2)\cdot e_2\cdot B_{12}\\ &-e_1\cdot j(e_2)\cdot B_{11}-e_2\cdot j(e_2)\cdot B_{12}+j(e_1)\cdot e_1\cdot B_{12}+j(e_1)\cdot e_2\cdot B_{22}\Big)\cdot\overline{\varphi}\\ =\;& \frac{i}{4}\Big(-2g(j(e_1),e_2)B_{22}+2g(j(e_2),e_1)B_{11}\Big)\cdot\overline{\varphi}\\ =\;& -i\,j_{12}H\cdot\overline{\varphi}. \qquad (39)\end{aligned}$$

Now, replacing (36) and (37) in (35) and combining together with (38) and (39), we get the desired result.

(4) Since j is antisymmetric, we have $j_{kl} = -j_{lk}$ and so

$$\begin{aligned}\mathcal{A}_6^{12} - \mathcal{A}_6^{21} &= \frac{1}{4}\big(j(e_2)\cdot j(e_1) - j(e_1)\cdot j(e_2)\big)\cdot\varphi \\ &= \frac{1}{2}j_{21}j_{12}e_1\cdot e_2\cdot\varphi = -\frac{1}{2}g(j(e_1),e_2)^2 e_1\cdot e_2\cdot\varphi \end{aligned} \tag{40}$$

(5)

$$\begin{aligned}&\mathcal{A}_7^{12} - \mathcal{A}_7^{21} \\ =\ & \frac{1}{4}(h(e_2)\cdot h(e_1) - h(e_1)\cdot h(e_2)) \\ =\ & \frac{1}{4}\big(-h_{21}h_{11} + h_{21}h_{12}\nu_1\cdot\nu_2 + h_{22}h_{11}\nu_2\cdot\nu_1 - h_{22}h_{12} + h_{11}h_{21} \\ & -h_{11}h_{22}\nu_1\cdot\nu_2 - h_{12}h_{21} - h_{11}h_{22}\nu_1\cdot\nu_2 - h_{12}h_{21}\nu_2\cdot\nu_1 + h_{12}h_{22}\big)\cdot\varphi \\ =\ & \frac{1}{2}(h_{21}h_{12} - h_{11}h_{22})\nu_1\cdot\nu_2\cdot\varphi\end{aligned}$$

(6)

$$\begin{aligned}\mathcal{A}_4^{12} - \mathcal{A}_4^{21} &= -\frac{i}{4}\big(e_2\cdot j(e_1) - j(e_2)\cdot e_1 - e_1\cdot j(e_2) + j(e_1)\cdot e_2\big)\cdot\overline{\varphi} \\ &= -\frac{i}{4}\big(-j_{12} + j_{21} + j_{21} - j_{12}\big)\cdot\overline{\varphi} = i\, j_{12}\overline{\varphi}.\end{aligned} \tag{41}$$

(7) We have

$$\begin{aligned}&\mathcal{A}_8^{12} - \mathcal{A}_8^{21} \\ =\ & \frac{1}{4}\big(j(e_2)\cdot h(e_1) + h(e_2)\cdot j(e_1) - j(e_1)\cdot h(e_2) - h(e_1)\cdot j(e_2)\big)\cdot\varphi \\ =\ & \frac{1}{2}\big(j_{21}h_{11}e_1\cdot\nu_1 + j_{21}h_{12}e_1\cdot\nu_2 + j_{21}h_{21}e_2\cdot\nu_1 + j_{12}h_{22}e_2\cdot\nu_2\big)\cdot\varphi\end{aligned} \tag{42}$$

Finally, we recall here that Lemma 3.3 of [4] says that first

$$d\eta(X,Y) = -\frac{1}{2}\sum_{j=1}^{2} e_j\cdot\big((\nabla'_X B)(Y,e_j)) - (\nabla'_Y B)(X,e_j)\big),$$

and second

$$\begin{aligned}\eta(e_2)\cdot\eta(e_1) - \eta(e_1)\cdot\eta(e_2) &= \frac{1}{2}\big(|B_{12}|^2 - \langle B_{11}, B_{22}\rangle\big)e_1\cdot e_2 \\ &\quad + \frac{1}{2}\langle(S_{\nu_1}\circ S_{\nu_2} - S_{\nu_2}\circ S_{\nu_1})(e_1), e_2\rangle\,\nu_1\cdot\nu_2.\end{aligned}$$

Moreover, since $d\eta(e_1,e_2) = \nabla_{e_1}(\eta(e_2)) - \nabla_{e_2}(\eta(e_1))$, we deduce immediately from the definition of $\mathcal{A}_1^{12}$ and $\mathcal{A}_1^{21}$ and the two above identities the desired relation.

□

Now, we have all the ingredients to prove Theorems 1.1 and 1.2.

5. Lagrangian case, proof of Theorem 1.2

First and from Sections 3 and 4, assertion (2) of Theorem 1.2 implies assertion (1). Assume now that assertion (1) is satisfied. Since, $j = t = 0$, we have

$$\mathcal{R}_{e_1,e_2}\varphi = \frac{1}{2}K_M e_1\cdot e_2\cdot\varphi - \frac{1}{2}K_E\nu_1\cdot\nu_2\cdot\varphi + \frac{1}{2}F^{M+E}(e_1,e_2)\varphi, \tag{43}$$

with $F^{M+E}(e_1,e_2) = 0$ because $j = 0$. On the other hand, we have

$$\begin{aligned}\mathcal{R}_{e_1,e_2}\varphi &= -\frac{1}{2}\sum_{j=1}^{2} e_j\cdot((\nabla^{'}_{e_1}B)(e_2,e_j)-(\nabla^{'}_{e_2}B)(e_1,e_j))\cdot\varphi\\ &\quad+\frac{1}{2}(|B_{12}|^2-<B_{11},B_{22}>)e_1\cdot e_2\cdot\varphi\\ &\quad+\frac{1}{2}<[S_{\nu_1},S_{\nu_2}](e_1),e_2>\nu_1\cdot\nu_2\cdot\varphi\\ &\quad-\frac{1}{2}e_1\cdot e_2\cdot\varphi+\frac{1}{2}(h_{21}h_{12}-h_{11}h_{22})\nu_1\cdot\nu_2\cdot\varphi \qquad (44)\end{aligned}$$

We get finally that $T\cdot\varphi=0$, where $T\in(\Lambda^2M\otimes 1\oplus TM\otimes E\oplus 1\otimes\Lambda^2E)$ is given by

$$\begin{aligned}T &= \frac{1}{2}(<B_{11},B_{22}>-|B_{12}|^2+1-K_M)e_1\wedge e_2\\ &\quad+\frac{1}{2}(h_{22}h_{11}-h_{21}h_{12}-<[S_{\nu_1},S_{\nu_2}](e_1),e_2>-K_E)\nu_1\wedge\nu_2\\ &\quad-\frac{1}{2}\sum_{j=1}^{2}e_j\wedge((\nabla^{'}_{e_1}B)(e_2,e_j)-(\nabla^{'}_{e_2}B)(e_1,e_j)).\end{aligned}$$

Now, we recall that Lemma 3.4 of [4] ensures that if T is a two form and φ a spinor so that φ^+ and φ^- never vanish and $T\cdot\varphi=0$, then $T=0$. Note that the hypothesis that both φ^+ and φ^- do not vanish is crucial. Here, the conclusion $T=0$ reduces to the following identities

$$K_M=<B_{11},B_{22}>-|B_{12}|^2+1,$$

$$K_E=-<[S_{\nu_1},S_{\nu_2}](e_1),e_2>-(h_{21}h_{12}-h_{22}h_{11}),$$

$$(\nabla^{'}_{e_1}B)(e_2,e_j)-(\nabla^{'}_{e_2}B)(e_1,e_j)=0,$$

which are Gauss, Ricci and Codazzi equations for a Lagrangian surface in $\mathbb{C}P^2$ and so the conditions (22) are fulfilled. Hence, by Proposition 3.2, we conclude that there exists a Lagrangian isomertic immersion from (M,g) into $\mathbb{C}P^2$ with E as normal bundle and B as second fundamental form. This proves that assertion (1) of Theorem 1.2 implies assertion (2). Theorem 1.2 is proved.

6. Complex case, proof of Theorem 1.1

Again, assertion (2) of Theorem 1.1 implies assertion (1) by the discussions of Sections 3 and 4. Assume now that assertion (1) is satisfied. We have $s=0$, $h=0$ so $F^{M+E}(e_1,e_2)=-2i$. We take $j(e_1)=e_2$ and $t\nu_1=\nu_2$, *i.e.* $g(j(e_1),e_2)=g(t\nu_1,\nu_2)=1$. We calculate and we get

$$\begin{aligned}\mathcal{R}_{e_1,e_2}\varphi &= -\frac{1}{2}K_Me_1\cdot e_2\cdot\varphi-\frac{1}{2}K_N\nu_1\cdot\nu_2\cdot\varphi+\frac{1}{2}F^{M+E}(e_1,e_2)\varphi\\ &= -\frac{1}{2}K_Me_1\cdot e_2\cdot\varphi-\frac{1}{2}K_N\nu_1\cdot\nu_2\cdot\varphi-i\varphi\\ &= \overline{T}\cdot\varphi-i\varphi, \qquad (45)\end{aligned}$$

where $\overline{T}$ is the 2-form defined by

$$\overline{T} = -\frac{1}{2}K_Me_1\wedge e_2-\frac{1}{2}K_N\nu_1\wedge\nu_2$$

On the other hand, we have

$$\begin{aligned}\mathcal{R}_{e_1,e_2}\varphi &= -e_1\cdot e_2\cdot\varphi+i\overline{\varphi}\\ &\quad -\frac{1}{2}\sum_{j=1}^{2}e_j\cdot((\nabla'_{e_1}B)(e_2,e_j)-(\nabla'_{e_2}B)(e_1,e_j))\cdot\varphi\\ &\quad +\frac{1}{2}(|B_{12}|^2-<B_{11},B_{22}>)e_1\cdot e_2\cdot\varphi\\ &\quad +\frac{1}{2}<[S_{\nu_1},S_{\nu_2}](e_1),e_2>\nu_1\cdot\nu_2\cdot\varphi\\ &\quad =\widetilde{T}\cdot\varphi+i\overline{\varphi},\end{aligned}\tag{46}$$

where $\widetilde{T}$ is the 2-form defined by

$$\begin{aligned}\widetilde{T} &= \frac{1}{2}(<B_{11},B_{22}>-|B_{12}|^2+1-K_M)e_1\wedge e_2\\ &\quad +\frac{1}{2}(h_{22}h_{11}-h_{21}h_{12}-<[S_{\nu_1},S_{\nu_2}](e_1),e_2>-K_E)\nu_1\wedge\nu_2\\ &\quad -\frac{1}{2}\sum_{j=1}^{2}e_j\wedge((\nabla'_{e_1}B)(e_2,e_j)-(\nabla'_{e_2}B)(e_1,e_j)).\end{aligned}$$

Together, it gives $\overline{T}\cdot\varphi-\widetilde{T}\cdot\overline{\varphi}-i\varphi-i\overline{\varphi}=0$, which means that $\mathcal{T}\cdot\varphi-i\varphi-i\overline{\varphi}=0$, where the 2-form $\mathcal{T}=\overline{T}-\widetilde{T}$ is given by

$$\begin{aligned}\mathcal{T} &= -\frac{1}{2}K_Me_1\wedge e_2\cdot\varphi-\frac{1}{2}K_N\nu_1\wedge\nu_2\cdot\varphi+e_1\wedge e_2\cdot\varphi\\ &\quad -\frac{1}{2}(|B_{12}|^2-<B_{11},B_{22}>)e_1\wedge e_2\cdot\varphi\\ &\quad -\frac{1}{2}<[S_{\nu_1},S_{\nu_2}](e_1),e_2>\nu_1\wedge\nu_2\cdot\varphi\\ &\quad +\frac{1}{2}\sum_{j=1}^{2}e_j\wedge((\nabla'_{e_1}B)(e_2,e_j)-(\nabla'_{e_2}B)(e_1,e_j))\cdot\varphi\end{aligned}\tag{47}$$

We give now the following Lemma:

Lemma 6.1. *Let $\mathcal{T}$ be a* 2 *form,* i.e. $\mathcal{T}\in\Lambda^2M\otimes 1\oplus\Lambda^1M\otimes\Lambda^1E\oplus 1\otimes\Lambda^2E$ *and $\varphi\in\Sigma$ so that both φ^+ and φ^- never vanish. Assume that*

$$\mathcal{T}\cdot\varphi-i\varphi-i\overline{\varphi}=0,$$

and write $\mathcal{T}=T^te_1\wedge e_2+T^n\nu_1\wedge\nu_2+T^m$, where $T^m\in\Lambda^1M\otimes\Lambda^1E$. Then,

$$T^t=-1,T^n=0 \text{ and } T^m=0.$$

Proof. Let $\varphi=\varphi^++\varphi^-$, with

$$\begin{aligned}\varphi^+&=\varphi^{++}+\varphi^{--},\\ \varphi^-&=\varphi^{-+}+\varphi^{+-},\end{aligned}$$

a solution of (23) with $h=0$. This means that

$$\begin{aligned}\nabla_X\varphi^{++}&=-\frac{1}{2}X\cdot\varphi^{-+}-\frac{i}{2}j(X)\cdot\varphi^{-+}\\ \nabla_X\varphi^{+-}&=-\frac{1}{2}X\cdot\varphi^{--}+\frac{i}{2}j(X)\cdot\varphi^{--}\\ \nabla_X\varphi^{-+}&=-\frac{1}{2}X\cdot\varphi^{++}+\frac{i}{2}j(X)\cdot\varphi^{++}\\ \nabla_X\varphi^{--}&=-\frac{1}{2}X\cdot\varphi^{+-}-\frac{i}{2}j(X)\cdot\varphi^{+-}.\end{aligned}$$

For a sake of simplicity, and without lost of generality, we can restrict only $\varphi^+ = \varphi^{++}$ and $\varphi^- = \varphi^{-+}$ which which have no zeros by assumption. The equation

$$\mathcal{T} \cdot \varphi - i\varphi - i\overline{\varphi} = 0,$$

becomes

$$T^t e_1 \cdot e_2 \cdot (\varphi^{++} + \varphi^{-+}) + (T^n + 1)\nu_1 \cdot \nu_2 \cdot (\varphi^{++} + \varphi^{-+}) + T^m \cdot (\varphi^{++} + \varphi^{-+}) = i\overline{\varphi} = i(\varphi^{++} - \varphi^{-+})$$

Taking the scalar product with φ^{++} then with φ^{-+}, we get

$$T^t + T^n + 1 = -1,$$
$$-T^t + T^n + 1 = 1,$$

which gives $T^n = -1$, $T^t = -1$ and $T^m = 0$. One can uses (47) to get Gauss, Codazzi and Ricci equations and so the conditions (21) are fulfilled. There are exactly the conditions of a complex immersion. Hence, by Proposition 3.2, we conclude that there exists a complex isometric immersion from (M, g) into $\mathbb{C}P^2$ with E as normal bundle and B as second fundamental form. As for the Lagrangian case, this proves that assertion (2) of Theorem 1.1 implies assertion (1).

7. The Dirac equation

Let φ be a spinor field satisfying Equation (23), then it satisfies the following Dirac equation

$$D\varphi = \vec{H} \cdot \varphi - \varphi + \frac{i}{2}\beta \cdot \overline{\varphi}, \tag{48}$$

where β is defined by $\beta = \sum_{i=1,2} e_i \cdot h(e_i) = \sum_{i,j=1}^{2} h_{ij} e_i \cdot \nu_j$, where $\{e_1, e_2\}$ and $\{\nu_1, \nu_2\}$ are respectively orthonormal frames of TM and E and $h_{ij} = \langle h(e_i), \nu_j \rangle$.
As in [4] and [35], we will show that this equation with an appropiate condition on the norm of both φ^+ and φ^- is equivalent to Equation (23), where the tensor B is expressed in terms of the spinor field φ and such that $\mathrm{tr}(B) = 2\vec{H}$. Moreover, from

$$\nabla_X \varphi^{\pm} = -\frac{1}{2}\eta(X) \cdot \varphi^{\pm} - \frac{1}{2} X \cdot \varphi^{\mp} \mp \frac{i}{2} j(X)\varphi^{\mp} \mp \frac{i}{2} h(X)\varphi^{\mp},$$

we deduce that

$$X(|\varphi^{\pm}|^2) = \mathrm{Re}\left\langle -\frac{1}{2} X \cdot \varphi^{\mp} \mp \frac{i}{2} j(X) \cdot \varphi^{\mp} \mp \frac{i}{2} h(X) \cdot \varphi^{\mp}, \varphi^{\pm} \right\rangle. \tag{49}$$

Now, let φ a spinor field solution of the Dirac equation (48) with φ^+ and φ^- nowhere vanishing and satisfying the norm condition (49), we set for any vector fields X and Y tangent to M and $\xi \in \Gamma(E)$

$$\begin{aligned} & \langle B(X,Y), \xi \rangle \\ = \; & \frac{1}{|\varphi^+|^2} \mathrm{Re}\left\langle X \cdot \nabla_Y \varphi^+ - \frac{1}{2}\left(X + ij(X) + ih(X)\right) \cdot Y \cdot \varphi^-, \xi \cdot \varphi^+ \right\rangle \\ & + \frac{1}{|\varphi^-|^2} \mathrm{Re}\left\langle X \cdot \nabla_Y \varphi^- - \frac{1}{2}\left(X - ij(X) - ih(X)\right) \cdot Y \cdot \varphi^-, \xi \cdot \varphi^+ \right\rangle \end{aligned} \tag{50}$$

Then, we have the following

Proposition 7.1. *Let $\varphi \in \Gamma(\Sigma)$ satisfying the Dirac equation*

$$D\varphi = \vec{H} \cdot \varphi - \varphi + \frac{i}{2}\beta \cdot \overline{\varphi}$$

such that

$$X(|\varphi^{\pm}|^2) = \mathrm{Re}\left\langle -\frac{1}{2} X \cdot \varphi^{\mp} \mp \frac{i}{2} j(X) \cdot \varphi^{\mp} \mp \frac{i}{2} h(X) \cdot \varphi^{\mp}, \varphi^{\pm} \right\rangle,$$

then φ is solution of Equation (23)

$$\nabla_X \varphi = -\frac{1}{2}\eta(X)\cdot\varphi - \frac{1}{2}X\cdot\varphi + \frac{i}{2}j(X)\cdot\overline{\varphi} + \frac{i}{2}h(X)\cdot\overline{\varphi},$$

where η is defined by $\eta(X) = \sum_{j=1}^{2} e_j \cdot B(e_j, X)$. Moreover, B is symmetric.

The proof of this proposition will not be given, since it is completely similar to the case of Riemannian products [35, Proposition 4.1]. Now, combining this proposition with Theorems 1.1 and 1.2, we get the following corollaries. We have this first one for complex immersions of surfaces.

Corollary 7.2. *Let (M^2, g) be an oriented Riemannian surface and E an oriented vector bundle of rank 2 over M with scalar product $<\cdot,\cdot>_E$ and compatible connection ∇^E. We denote by $\Sigma = \Sigma M \otimes \Sigma E$ the twisted spinor bundle. Let j be a complex structure on M and t a complex structure on E. Let $\vec{H}$ be a section of E. Then, the following two statements are equivalent*

(1) *There exists a Spin^c structure on $\Sigma M \otimes \Sigma E$ whose auxiliary line bundle's curvature is given by $F^{M+E}(e_1, e_2) = 0$ and a spinor field φ in Σ solution of the Dirac equation*

$$D\varphi = \vec{H}\cdot\varphi - \varphi$$

such that φ^+ and φ^- never vanish satisfying the norm condition

$$X(|\varphi^{\pm}|^2) = \mathrm{Re}\left\langle -\frac{1}{2}X\cdot\varphi^{\mp} \mp \frac{i}{2}j(X)\cdot\varphi^{\mp}\varphi^{\pm}\right\rangle$$

and such that the maps j, t and the tensor B defined by (50) *satisfies $t(B(X,Y)) = B(X, j(Y))$ for all $X, Y \in \Gamma(TM)$.*

(2) *There exists an isometric* **complex** *immersion of (M^2, g) into $\mathbb{C}P^2$ with E as normal bundle and mean curvature $\vec{H}$ such that over M the complex structure of $\mathbb{C}P^2$ is given by j and t (in the sense of Proposition 3.2).*

We have this second corollary for Lagrangian surfaces.

Corollary 7.3. *Let (M^2, g) be an oriented Riemannian surface and E an oriented vector bundle of rank 2 over M with scalar product $<\cdot,\cdot>_E$ and compatible connection ∇^E. We denote by $\Sigma = \Sigma M \otimes \Sigma E$ the twisted spinor bundle. Let $B : TM \times TM \longrightarrow E$ a bilinear symmetric map, $h : TM \longrightarrow E$ and $s : E \longrightarrow TM$ the dual map of h. Assume that the maps h, s are parallel and satisfy $h \circ s = -\mathrm{id}_E$. Let $\vec{H}$ be a section of E. Then, the following two statements are equivalent*

(1) *There exists a Spin^c structure on $\Sigma M \otimes \Sigma E$ whose auxiliary line bundle's curvature is given by $F^{M+E}(e_1, e_2) = -2i$ and a spinor field φ in Σ solution of the Dirac equation*

$$D\varphi = \vec{H}\cdot\varphi - \varphi + \frac{i}{2}\beta\cdot\overline{\varphi}$$

$\big(\beta$ is the 2-form defined by $\beta = \sum_{i=1,2} e_i \cdot h(e_i)\big)$ such that φ^+ and φ^- never vanish satisfying the norm condition

$$X(|\varphi^{\pm}|^2) = \mathrm{Re}\left\langle -\frac{1}{2}X\cdot\varphi^{\mp} \mp \frac{i}{2}h(X)\cdot\varphi^{\mp}\varphi^{\pm}\right\rangle$$

and such that the tensor B defined by (50) *satisfies $A_{h(Y)}X + s(B(X,Y)) = 0$, for all $X \in TM$, where $A_\nu : TM \longrightarrow TM$ if defined by $g(A_\nu X, Y) = \langle B(X,Y), \nu\rangle_E$ for all $X, Y \in \Gamma(TM)$ and $\nu \in \Gamma(E)$.*

(2) *There exists an isometric* **Lagrangian** *immersion of* (M^2, g) *into* $\mathbb{C}P^2$ *with* E *as normal bundle and mean curvature* $\vec{H}$ *such that over* M *the complex structure of* $\mathbb{C}P^2$ *is given by* h *and* s *(in the sense of Proposition 3.2).*

Acknowledgment. The first named author would like to thank the University of Paris-Est, Marne La Vallée for its support and hospitality. The authors warmly thank the referee for his/her remraks that allow to highly improve the present paper. Both authors are also grateful to Mihaela Pilca for helpful discussions about Kählerian Killing spinors.

REFERENCES

[1] C. BÄR, *Extrinsic bounds for eigenvalues of the Dirac operator*, Ann. Glob. Anal. Geom. 16 (1998) 573-596.
[2] W. BALLMANN,l *Lectures on Kähler manifolds*, ESI Lectures in Mathematics and Physics, ISBN 978-3-03719-025-8, 2006.
[3] P. BAYARD, *On the spinorial representation of spacelike surfaces into 4-dimensional Minkowski space*, J. Geom. Phys. 74 (2013), 289-313.
[4] P. BAYARD, M.A. LAWN & J. ROTH, *Spinorial representation of surfaces in four-dimensional Space Forms*, Ann. Glob. Anal. Geom. 44 (4) (2013), 433-453.
[5] P. BAYARD & V. PATTY, *Spinor representation of Lorentzian surfaces in* $\mathbb{R}^{2,2}$, J. Geom. Phys. 95 (2015), 74-95.
[6] A. L. BESSE, *Einstein Manifolds*, Ergebnisse der Mathematik (3), Springer Verlag, Berlin (1987).
[7] J. P. BOURGUIGNON, O. HIJAZI, J. L. MILHORAT, A. MOROIANU & S. MOROIANU, *A spinorial approach to Riemannian and conformal geometry*, EMS Monographs in Mathematics, 2015.
[8] B. Daniel, *Isometric immersions into* $\mathbb{S}^n \times \mathbb{R}$ *and* $\mathbb{H}^n \times \mathbb{R}$ *and applications to minimal surfaces*, Trans. Amer. Math. Soc. 361 (2009) no 12, 6255-6282.
[9] B. Daniel, *Isometric immersions into* 3*-dimensional homogeneous manifolds*, Comment. Math. Helv. 82 (2007), 87-131.
[10] T. FRIEDRICH, *On the spinor representation of surfaces in Euclidean* 3*-space*, J. Geom. Phys. (1998), 143-157.
[11] N. GINOUX, *Opérateurs de Dirac sur les sous-variétés*, Ph. D thesis 2002, Institut Élie Cartan, Nancy.
[12] N. GINOUX & B. MOREL, *On eigenvalue estimates for the submanifold Dirac operator*, Int. J. Math. 13 (2002), No. 5, 533-548.
[13] N. GINOUX, *The Dirac spectrum*, Lect. Notes in Math. 1976, Springer 2009.
[14] F. HÉLEIN & P. ROMON, *Weierstrass representation of Lagrangian surfaces in four-dimensional space using spinors and quaternions*, Comment. Math. Helv. 75 (2000), 668-680.
[15] F. HÉLEIN & P. ROMON, *Hamiltonian stationary tori in the complex projective plane*, Proc. London Math. Soc. 90 (2005), 472-496.
[16] O. HIJAZI, S. MONTIEL & F. URBANO, Spin^c *geometry of Kähler manifolds and the Hodge Laplacian on minimal Lagrangian submanifolds*, Math. Z. 253, Number 4 (2006) 821-853.
[17] O. HIJAZI, *Spectral properties of the Dirac operator and geometrical structures*, Proceedings of the summer school on geometric methods in quantum field theory, Villa de Leyva, Colombia, July 12-30, 1999, World Scientific 2001.
[18] N. HITCHIN, *Harmonic spinors*, Adv. Math.14 (1974), 1-55.
[19] K.-D. KIRCHBERG, *An estimation for the first eigenvalue of the Dirac operator on closed Kähler manifolds of positive scalar curvature*, Ann. Global Anal. Geom. 4 (1986), 291-325.
[20] B.G. KONOPELCHENKO, *Weierstrass representations for surfaces in 4D spaces and their integrable deformations via DS hierarchy*, Ann. Glob. Anal. Geom. 18 (2000) 61-74.
[21] D. KOWALCZYK, *Isometric immersions into products of space forms*, Geom. Dedicata 151 (2011), 1-8.
[22] R. KUSNER AND N. SCHMIDT, *The spinor representation of surfaces in space*, Preprint arxiv dg-ga/9610005 (1996).
[23] M.A. LAWN, *A spinorial representation for Lorentzian surfaces in* $\mathbb{R}^{2,1}$, J. Geom. Phys. 58 (2008) no. 6, 683-700.
[24] M.-A. LAWN & M. ORTEGA, *A Fundamental Theorem for Hypersurfaces in Semi-Riemannian Warped Products*, J. Geom. Phys. 90 (2015), 55-70.
[25] M.A. LAWN & J. ROTH, *Spinorial characterization of surfaces in pseudo-Riemannian space forms*, Math. Phys. Anal. and Geom. 14 (2011) no. 3, 185-195.
[26] M.A. LAWN & J. ROTH, *Isometric immersions of hypersurfaces into 4-dimensional manifolds via spinors*, Diff. Geom. Appl. 28 (2) (2010), 205-219
[27] H. B. LAWSON & M.-L. MICHELSON, *Spin Geometry*, Princeton Mathematical Series, 38. Princeton University Press (1989).
[28] S. MONTIEL, *Using spinors to study submanifolds*, Roma 2004 - Nancy 2005.

[29] B. MOREL, *Surfaces in $\mathbb{S}^3$ and $\mathbb{H}^3$ via spinors*, Actes du séminaire de théorie spectrale, Institut Fourier, Grenoble, 23 (2005), 9-22.
[30] A. MOROIANU, *Lectures on Kähler Geometry*, London Mathematical Society Student Text 69, Cambridge University Press, Cambridge, 2007.
[31] A. MOROIANU, *Parallel and Killing spinors on Spin^c manifolds*, Comm. Math. Phys. 187 (1997), 417-427.
[32] R. NAKAD & J. ROTH, *Hypersurfaces of $Spin^c$ manifolds and Lawson Type correspondence*, Annals of Global Analysis and Geometry, Vol 42 no 3 (2012), pp 421-442.
[33] P. PICCIONE AND D.V. TAUSK, *An existence theorem for G-structure preserving affine immersions*, Indiana Univ. Math. J. 57 (3) (2008), 1431-1465.
[34] P. ROMON & J. ROTH, *The spinor representation formula in 3 and 4 dimensions*, Pure and Applied Differential Geometry, Proceedings of the conference PADGE 2012, Shaker Verlag, Aachen (2013) 261-282.
[35] J. ROTH, *Spinorial characterization of surfaces into 3-dimensional homogeneous manifolds*, J. Geom. Phys 60 (2010), 1045-106.
[36] J. ROTH, *Isometric immersion into Lorentzian products*, Int. J. Geom. Method. Mod. Phys, 8 (2011) no.6, 1-22.
[37] J. ROTH, *Spinors and isometric immersions of surfaces into 4-dimensional products*, Bulletin of the Belgian Mathematical Society - Simon Stevin Vol 21 no 4 (2014), pp 635-652.
[38] I. TAIMANOV, *Surfaces of revolution in terms of solitons*, Ann. Glob. Anal. Geom. 15 (1997) 410-435.
[39] I. TAIMANOV, *Surfaces in the four-space and the Davey-Stewartson equations*, J. Geom Phys. 56 (2006) 1235-1256.
[40] F. URBANO, *Hamiltonian stability and index of minimal Lagrangian surfaces of the complex projective plan*, Indiana Univ. Math. J. 56 (2007), 931-946.

(R. Nakad) NOTRE DAME UNIVERSITY-LOUAIZÉ, FACULTY OF NATURAL AND APPLIED SCIENCES, DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O. BOX 72, ZOUK MIKAEL, LEBANON,

E-mail address: rnakad@ndu.edu.lb

(J. Roth) LAMA, UNIVERSITÉ PARIS-EST MARNE-LA-VALLÉE, CITÉ DESCARTES, CHAMPS SUR MARNE, 77454 MARNE-LA-VALLÉE CEDEX 2, FRANCE

E-mail address: julien.roth@u-pem.fr